

Available at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/IJMYCO

Performance comparison of a pair of Lowenstein–Jensen media supplemented with pyruvate or glycerol, and the combination of both supplements in a single Lowenstein–Jensen medium for the growth support of the *Mycobacterium Tuberculosis* complex

Aliou K. Faburay^{a,b,*}, Francis S. Mendy^a, Jacob K. Otu^a, Tutty Isatou Faal-Jawara^a, F. Gehre^{a,b}, Ousman Secka^a

^a Medical Research Council Unit, Serrekunda, the Gambia, Serrekunda, Gambia

^b Institute of Tropical Medicine, Antwerp, Belgium

ARTICLE INFO

Article history:

Received 21 September 2016

Accepted 23 September 2016

Available online xxxx

Keywords:

Lowenstein–Jensen media

Mycobacterium tuberculosis complex,

Lowenstein–Jensen (LJ)

Pyruvate

Glycerol

ABSTRACT

Objective/Background: To evaluate the performance of Lowenstein–Jensen medium (LJ) supplemented with pyruvate and glycerol (LJPG), compared with LJ supplemented with pyruvate (LJP) or glycerol (LJG) for the support of mycobacterial growth.

Method: This study used 100 Ziehl–Neelsen-confirmed positive mycobacterium growth indicator tube 960 culture samples that were obtained from clinical samples during routine diagnosis. All cultures were inoculated in parallel on LJG/LJP and on LJGP, which were incubated and read weekly for the evidence of growth. The mycobacterial recovery rate, contamination rate, and time to detection were compared.

Result: The recovery rate for LJG/LJP and for LJPG was 90% (90 samples) and 88% (88 samples), respectively (kappa p-value, 0.9). There was no significant difference in the contamination rate, which was 8% (8 samples) for LJG/LJP and 9% (9 samples) for LJPG. Mycobacterial growth was faster in LJPG (1.6 weeks) than in LJG/LJP (2 weeks).

Conclusion: A single LJPG slope was not significantly different, compared with the usual pair of LJG or LJP slopes. This is a promising new culturing approach that could be used in *Mycobacterium africanum*-endemic in West African countries. It significantly reduces labor time and consumable costs and more quickly detects the *M. tuberculosis* complex.

Conflicts of interest

The authors have no conflicts of interest to declare.

* Corresponding author.

E-mail address: faburayak@hotmail.com (A.K. Faburay).

Peer review under responsibility of Asian African Society for Mycobacteriology.

<http://dx.doi.org/10.1016/j.ijmyco.2016.09.019>